

This is the published version of a paper published in *Environmental Research Letters*.

Citation for the original published paper (version of record):

Gustavsson, L., Sathre, R., Leskinen, P., Nabuurs, G-J., Kraxner, F. (2022) Comment on 'Climate mitigation forestry—temporal trade-offs' *Environmental Research Letters*, 17(4): 048001 https://doi.org/10.1088/1748-9326/ac57e3

Access to the published version may require subscription.

N.B. When citing this work, cite the original published paper.

Permanent link to this version:

http://urn.kb.se/resolve?urn=urn:nbn:se:lnu:diva-110895

ENVIRONMENTAL RESEARCH

LETTERS

COMMENT • OPEN ACCESS

Comment on 'Climate mitigation forestry—temporal trade-offs'

To cite this article: Leif Gustavsson et al 2022 Environ. Res. Lett. 17 048001

View the <u>article online</u> for updates and enhancements.

You may also like

- Can life-cycle assessment produce reliable policy quidelines in the building sector?
 Antti Säynäjoki, Jukka Heinonen, Seppo Junnila et al.
- Phonon-mediated back-action of a charge readout on a double quantum dot U Gasser, S Gustavsson, B Küng et al.
- Comparative Life-Cycle Analysis of Building Materials for the Thermal Upgrade of an Existing Building C Piccardo, A Dodoo, L Gustavsson et al.

ENVIRONMENTAL RESEARCH

LETTERS



OPEN ACCESS

RECEIVED

22 November 2021

REVISED

3 January 2022

ACCEPTED FOR PUBLICATION 23 February 2022

11 March 2022

Original content from this work may be used under the terms of the Attribution 4.0 licence.

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



COMMENT

Comment on 'Climate mitigation forestry—temporal trade-offs'

Leif Gustavsson^{1,*}, Roger Sathre¹, Pekka Leskinen², Gert-Jan Nabuurs³ and Florian Kraxner

- Linnaeus University, Växjö, Sweden
- European Forest Institute, Joensuu, Finland
- Wageningen University and Research, Wageningen, The Netherlands
- International Institute for Applied Systems Analysis, Laxenburg, Austria
- Author to whom any correspondence should be addressed.

E-mail: leif.gustavsson@lnu.se

In an article recently published in ERL, Skytt et al (2021) describe a modeling exercise in five Swedish counties that found short-term climate benefits from reducing forest harvest. We agree with Skytt et al that forests have strong potential for climate change mitigation, that effects and consequences must be considered over the short-, medium-, and long-terms, and that old forests are important and fulfil many functions including biodiversity protection. However, we argue that Skytt et al (a) place unfounded faith in the ability to quickly develop and deploy sustainable non-forest-based supply chains to respond to reduced forest harvest and reduced supply of wood-based materials and energy, (b) violate life cycle assessment (LCA) standards by not maintaining a consistent functional unit, (c) apply substitution factors (SF) for forest-based energy and materials that are low compared to current scientific evidence, (d) do not consider the climate mitigation potential of harvesting sustainable shares of forest residues, and (e) do not consider the effects of climate change and the risk of disturbance to carbon stored in forest ecosystems.

(a) We know that mitigation of, and adaptation to, climate change requires a strategic evolution and transformation of various technical sectors including energy, manufacturing, construction, and transport (Johnsson et al 2019, Tong et al 2019, Cowie at al 2021, UNEP 2021). For example, massive deployment of high efficiency renewable energy systems is needed. Fully avoiding climate disruption will increasingly require negative emission technologies, such as bioenergy with carbon capture and storage. Restricting forest products will limit the options available for strategic evolution and transformation of energy and material sectors and will make the implementation of sustainable technical systems more challenging. Skytt et al focus on shortand medium term carbon stock benefits from

reduced harvest. They argue that short term gains in forest carbon stock will 'buy us the time needed to implement sustainable technical systems, such as non-fossil fuel based electricity production, carbon capture and storage etc.' Yet they assume a SF of zero for electricity in the modelling exercise, implying that such systems have already been fully deployed. They also offer an 'illustrative example' of using hydrogen for steelmaking, an immature technology still at the research stage and with uncertain prospects. Skytt et al appear to underestimate the challenge of rapidly transitioning from our current dependence on non-renewable energy and material resources.

(b) Skytt et al do not consider the decreased supply of pulp and paper products to end users when forest harvest levels and the production of pulp and paper are decreased in their study. This is a severe methodological shortcoming, which violates LCA standards by fulfilling different functional units when comparing different forest management alternatives. The LCA methodology framed by ISO14040 and 14044 standards (ISO 2006a, 2006b) is intended to compare alternatives on a functionally equivalent basis. Skytt et al fail to ensure functional equivalency by ignoring the services provided by pulp and paper products, which would decrease in step with decreased forest harvest. They justify this omission with reference to Sathre and O'Connor (2010) and Leskinen et al (2018), saying they 'included only cases where a decrease of the supply of biomass to the industry would lead to increased use of fossil fuels or materials.' This is a misuse of the recommendations of Sathre and O'Connor (2010) and Leskinen et al (2018), which are only applicable to calculating substitution benefits in greenhouse gas balances, and are no justification for violating LCA standards by failing to fulfill a consistent

- functional unit. Skytt *et al* then call out 'the need to investigate to what extent different product groups provide substitution. It is only in cases where reduced use of wood products lead (sic) to increased future use of fossil-based products or fuels, that avoided emissions can be accounted.' Yet in the case of reducing forest harvest, it is clear that 100% of the reduced service once provided by forest products, will need to be substituted to maintain identical services and fulfil the functional unit as per LCA standards.
- (c) We agree with Skytt et al that SF can be a significant component when estimating the total mitigation potential of the forest sector. Skytt et al argue that technological development and renewable energy deployment will steadily improve the environmental performance of non-forest materials over time, thus reducing the climate SF of forest products. However, advancements are not limited to non-forest sectors, and current development within the forest sector offers improvements in forest management, wood-based materials, and forest bioenergy technology. Furthermore, given the urgency of climate change mitigation, the forest industry is actively seeking improved product portfolios to increase the climate benefits of wood-based products in replacing fossil materials and energy. Examples of this include wood-based textiles and modern wood construction materials such as cross-laminated timber (CLT). Novel technologies will typically develop faster than existing, more mature technologies. Furthermore, there will be substitution opportunities on the margin for a long time, as fossil fuels and carbon intensive materials remain in use in many areas. Therefore, material and energy substitution activities can be, and should be, targeted for high effect, thus specific marginal SF values should be used and not current average values. The SF values for material and energy used by Skytt et al are low compared to current scientific evidence. For example, Skytt et al used default SF values for energy that include a SF for electricity of zero. However, rough SF values for standalone electricity production including fuel cycle and end use emissions are 0.43 and 0.98 when woody biomass replaces fossil gas and coal, respectively⁵.
- ⁵ Calculations are based on fuel-to-electricity conversion efficiencies of 50% for fossil gas and 40% for coal and biomass, default fuel combustion emissions from IPCC (2006), and upstream fuel cycle fossil emissions of 5% of combustion emissions for all fuels. The corresponding SF values are 0.46 and 0.91 for state-of-theart energy technologies based on fuel-to-electricity conversion efficiencies of 56% for fossil gas, 46% for coal, and 45% for biomass (Danish Energy Agency 2020), and upstream and combustion emissions from IPCC (2006), Gustavsson *et al* (2015) and Gode *et al* (2011), and considering Swedish logging residues with an international transport of 1000 km. Using state-of-the-art

- Valid SF values are important in this sector, as woody residues from forests and forest industries can efficiently generate electricity in standalone and cogeneration power plants, in place of fossil fuels. We agree with Skytt et al that reduced use of wood products will reduce electricity use within the forest industry, but we note that electricity use will increase in other industries that make alternatives to wood products. Overall, the significance of accurate SF values is seen by comparing figures 7(a) and (b) of Skytt et al (2021): with higher SF values corresponding to modern forest product usage, the initial period during which reducing harvest may be climatically beneficial is shortened to mere decades, followed by strong climate impact thereafter.
- Skytt *et al* do not consider the harvest of logging residues in their forest management alternatives, despite such practice having a significant potential for renewable energy supply in Sweden. The annual current Swedish harvest of forest slash is about 10 TWh, while the annual potential slash and stump harvest may be about 65 and 40 TWh, respectively (IRENA 2019). Due to Swedish soil conditions a large extraction of logging residues could lead to a deficiency of nutrients and a reduction in forest productivity (Koponen et al 2015), thus sustainable management could require ash recycling and selective fertilization to ensure forest productivity. The Swedish Forest Agency (2019) has recommendations regarding the extraction of forest residues and the application of recycled ash to secure sustainable harvest levels while considering soil fertility and biodiversity. Replacing fossil fuels with harvested logging residues, which would otherwise decay naturally and release their stored carbon, will help to mitigate climate change (Sathre and Gustavsson 2011, Gustavsson et al 2017, 2021). Hence, the potential climate benefits are underestimated by Skytt et al, particularly for forest management alternatives with higher harvest levels.
- (e) Skytt *et al* briefly mention the 'effects of changes in precipitation and risks of damage from extreme weather events and pests', but their analysis does not consider potential disturbances to long-term carbon storage in forests. Climate change is not considered in their modeling, yet is expected to cause various impacts to Nordic forests, which are projected to benefit in terms of higher productivity but face higher risk of disturbances. Relying on indefinite carbon storage in forest ecosystems is a risky climate

technologies show rather small changes of SF compared to default values, as the technological development improves the conversion efficiencies of both the fossil and bioenergy technologies. However, more efficient technologies are important as more electricity is produced per unit of fuel.

mitigation strategy, particularly in the context of future temperature rise, precipitation variability, and altered disturbance regimes. Skytt *et al* ignore these risks, unrealistically assuming that unharvested forests will continue to store carbon indefinitely.

We agree with Skytt et al that science-based publications have reached different conclusions about forestry and its climate effects. The large global variation in forest ecology, forest management, and energy and material systems can partially explain these differences. However, different methodological approaches also explain the variation, especially between studies in the same ecological and technological context (Cowie et al 2021). Cowie et al conclude that focusing on short-term emissions reduction could make it more difficult to achieve medium- and long-term reductions, and that narrow perspectives obscure the most important role that bioenergy can play: to support the transformation of energy, industrial, and transport systems so that fossil fuels remain stored in geological formations.

Skytt et al are mainly focused on 'the coming 10-30 years' and they 'suggest that future research should focus on forest management strategies that can provide rapid climate benefits.' While appreciating the importance of the short term, we also must consider livelihood strategies in the medium- and long-term perspectives. Skytt et al favor the onetime climate benefit of increasing forest stock, at the expense of the continuous climate benefits of utilizing forest flows. Forest system modeling that is unencumbered by the flaws described above shows that reducing forest cuttings in south Sweden offers some climate benefits in the short term of 30-40 yr (Gustavsson et al 2021), in line with the results for all of Sweden (Gustavsson et al 2017). But after this initial period, the climate benefits are greater for active forestry with high harvest and efficient utilization of biomass. After 200 years such active forestry may generate about ten times greater carbon emission reduction, compared to the initial reduction from limiting forest cuttings (Gustavsson et al 2021). Many other studies also acknowledge that forest biomass as part of ongoing forest value chains can contribute to climate change mitigation, especially in the medium to long term (e.g. Marland and Schlamadinger 1997, Kraxner et al 2003, Lundmark et al 2014, Smyth et al 2014, Creutzig et al 2015, Kilpeläinen et al 2016, Favero et al 2017, 2020, Nabuurs et al 2017, Vance 2018, Petersson et al 2021). Hence, we argue that forests fulfil many functions, and in a country like Sweden the climate goals can best be met with active forestry including harvest and efficient utilization of renewable biomass for replacement of carbon-intensive non-wood products and fuels.

ORCID iDs

Roger Sathre https://orcid.org/0000-0002-1861-6423

Gert-Jan Nabuurs https://orcid.org/0000-0002-9761-074X

Florian Kraxner https://orcid.org/0000-0003-3832-6236

References

Cowie A L *et al* 2021 Applying a science-based systems perspective to dispel misconceptions about climate effects of forest bioenergy *Glob. Change Biol. Bioenergy* 13 1210–31

Creutzig F *et al* 2015 Bioenergy and climate change mitigation: an assessment *Glob. Change Biol. Bioenergy* 7 916–44

Danish Energy Agency 2020 Technology data: generation of electricity and district heating

Favero A, Daigneault A and Sohngen B 2020 Forests: carbon sequestration, biomass energy, or both? *Sci. Adv.* **6** eaay6792

Favero A, Mendelsohn R and Sohngen B 2017 Using forests for climate mitigation: sequester carbon or produce woody biomass? *Clim. Change* 144 195–206

Gode J, Martinsson F, Hagberg L, Öman A, Höglund J and Palm D 2011 Estimated Emission Factors for Fuels, Electricity, Heat and Transport in Sweden—Miljöfaktaboken 2011 (Stockholm: Värmeforsk)

Gustavsson L, Haus S, Lundblad M, Lundström A, Ortiz C A, Sathre R, Le Truong N and Wikberg P E 2017 Climate change effects of forestry and substitution of carbon-intensive materials and fossil fuels *Renew. Sustain. Energy Rev.* 67 612–24

Gustavsson L, Haus S, Ortiz C, Sathre R and Truong N L 2015 Climate effects of bioenergy from forest residues in comparison to fossil energy *Appl. Energy* **138** 36–50

Gustavsson L, Nguyen T, Sathre R and Tettey U Y A 2021 Climate effects of forestry and substitution of concrete buildings and fossil energy *Renew. Sustain. Energy Rev.* 136 110435

IPCC (Intergovernmental Panel on Climate Change) 2006 Guidelines for national greenhouse gas inventories

IRENA (International Renewable Energy Agency) 2019 Bioenergy from boreal forests: Swedish approach to sustainable wood use

ISO (International Organization for Standardization) 2006a ISO 14040: environmental management—life cycle assessment—principles and framework

ISO (International Organization for Standardization) 2006b ISO 14044: environmental management—life cycle assessment—requirements and guidelines

Johnsson F, Kjärstad J and Rootzén J 2019 The threat to climate change mitigation posed by the abundance of fossil fuels *Clim. Policy* **19** 258–74

Kilpeläinen A, Torssonen P, Strandman H, Kellomäki S, Asikainen A and Peltola H 2016 Net climate impacts of forest biomass production and utilization in managed boreal forests *Glob. Change Biol. Bioenergy* **8** 307–16

Koponen K *et al* 2015 Sustainability of forest energy in Northern Europe *VTT Technol.* **237** (https://publications.vtt.fi/ pdf/technology/2015/T237.pdf)

Kraxner F, Nilsson S and Obersteiner M 2003 Negative emissions from BioEnergy use, carbon capture and sequestration (BECS)—the case of biomass production by sustainable forest management from semi-natural temperate forests *Biomass Bioenergy* 24 285–96

Leskinen P, Cardellini G, González-García S, Hurmekoski E, Sathre R, Seppälä J, Smyth C, Stern T and Verkerk P J 2018 Substitution effects of wood-based products in climate change mitigation *From Science to Policy* 7 (European Forest Institute)

- Lundmark T, Bergh J, Hofer P, Lundström A, Nordin A, Poudel B C, Sathre R, Taverna R and Werner F 2014 Potential roles of Swedish forestry in the context of climate change mitigation *Forests* 5 557–78
- Marland G and Schlamadinger B 1997 Forests for carbon sequestration or fossil fuel substitution? A sensitivity analysis Biomass Bioenergy 13 389–97
- Nabuurs G J, Delacote P, Ellison D, Hanewinkel M, Hetemäki L and Lindner M 2017 By 2050 the mitigation effects of EU forests could nearly double through climate smart forestry Forests 8 484
- Petersson H, Ellison D, Mensah A A, Berndes G, Egnell G, Lundblad M, Lundmark T, Lundström A, Stendahl J and Wikberg P E 2021 On the role of forests and the forest sector for climate change mitigation in Sweden Manuscript Zenodo
- Sathre R and Gustavsson L 2011 Time-dependent climate benefits of using forest residues to substitute fossil fuels *Biomass Bioenergy* 35 2506–16
- Sathre R and O'Connor J 2010 Meta-analysis of greenhouse gas displacement factors of wood product substitution *Environ*. *Sci. Policy* 13 104–14

- Skytt T, Englund G and Jonsson B G 2021 Climate mitigation forestry—temporal trade-offs *Environ. Res. Lett.* **16** 114037
- Smyth C E, Stinson G, Neilson E, Lemprière T C, Hafer M, Rampley G J and Kurz W A 2014 Quantifying the biophysical climate change mitigation potential of Canada's forest sector *Biogeosciences* 11 3515–29
- Swedish Forest Agency 2019 Regler och rekommendationer för skogsbränsleuttag och kompensationsåtgärder (Rules and recommendations for forest fuel extraction and compensation measures) *Report 2019/14* (in Swedish)
- Tong D, Zhang Q, Zheng Y, Caldeira K, Shearer C, Hong C, Qin Y and Davis S J 2019 Committed emissions from existing energy infrastructure jeopardize 1.5 °C climate target Nature 572 373–7
- UNEP (United Nations Environment Programme) 2021 Making peace with nature: a scientific blueprint to tackle the climate, biodiversity and pollution emergencies
- Vance E D 2018 Conclusions and caveats from studies of managed forest carbon budgets *For. Ecol. Manage.* 427 350–4